

کزین برتر اندیشه برنگذرد
خداوند روزی ده رهنمای
فروزنده ماه و ناهید و مهر
نگارندهی بر شده پیکر است
نبینی مرنجان دو بیننده را
که او برتر از نام و از جایگاه
نیابد بدو نیز اندیشه راه
همان را گزیند که بیند همی
میان بندگی را ببایدست بست
در اندیشه‌ی سخته کی گنجد اوی
ستود آفریننده را کی توان
ز گفتار بی کار یکسو شوی
به ژرفی به فرمانش کردن نگاه
ز دانش دل پیر برنا بود
ز هستی مر اندیشه را راه نیست

به نام خداوند جان و خرد
خداوند نام و خداوند جای
خداوند کیوان و گردان سپهر
ز نام و نشان و گمان برترست
به بینندگان آفریننده را
نیابد بدو نیز اندیشه راه
سخن هر چه زین گوهران بگذرد
خرد گر سخن برگزیند همی
ستودن نداند کس او را چو هست
خرد را و جان را همی سنجد اوی
بدین آلت رای و جان و زبان
به هستیش باید که خستو شوی
پرستنده باشی و جوینده راه
توانا بود هر که دانا بود
از این پرده برتر سخنگاه نیست



Bu-Ali Sina University

Faculty of Chemistry

THESIS

Submitted in Partial Fulfillment of the Requirements for the Degree of
Doctor of Philosophy (Ph.D.) in INORGANIC CHEMISTRY

Title:

**Synthesis and characterization of macroacyclic
and macrocyclic complexes derived from various
mono- and di- aldehydes and some polyamines in
the presence of different metal ions**

Under Supervision of:

Professor Hassan Keypour

Advisor:

Professor S. Javad Sabounchei

By:

Reza Azadbakht

February 2008

۸۷/۱۰/۱۹۹۳
۸۷/۱۰/۲۴



دانشکده شیمی

پایان نامه برای دریافت درجه دکتری (Ph.D.) شیمی معدنی

عنوان :

سنتز و شناسایی کمپلکس های بزرگ حلقه و بزرگ غیر حلقه مشتق شده از دی آلدئیدها و منوآلدئیدهای مختلف و برخی از پلی آمینها در حضور تعدادی از یونهای فلزی

استاد راهنما :

پروفسور حسن کی پور

استاد مشاور :

پروفسور سید جواد صابونچی

پژوهشگر :

رضا آزادبخت

۱۳۸۷/۱۰/۱۳

بهمن ماه ۱۳۸۶

۱۰۷۷۳۸

All rights reserved. No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, recording, or otherwise, without the prior written permission of the Bu-Ali Sina university

Dedicated to:

My dear and honourable parents

*My dear and kind wife who accompanied
me all the time*

And

My dear and lovely daughter


Rozhan

Acknowledgment

This document of work could not been achieved without the help of those around me. I wish to take this opportunity to express my thanks to a number of people who have influenced this work:

- *First and foremost, my supervisor, Professor Hassan Keypour, for giving me the chance to undertake this project and for his guidance, patience, constant support, continual encouragement, and for many other things.*
- *Deep acknowledgments are offered to the honourable members of the thesis committee; Prof. S. J. Sabounchei; Prof. N. Safari; Prof. S. Amani; Prof. M.A. Zolfigol and Assoc. Prof. S. Salehzadeh.*
- *I wish to thank all members of the academic and technical staff who made my stay at the Bu-Ali Sina University useful and enjoyable experience. I learn many things from you and for this and many other things you have my thanks.*
- *A debt of gratitude is due to the other members of the Prof. H. Keypour research group. I thank you all for your friendship, assistance and cooperation over the last several years.*
- *Deep acknowledgments are offered to my dear and honourable parents.*
- *Very special thanks must be go to my precious wife, who have been an absolute pillar of strength. I appreciate your understanding, patience, love and willingness to sacrifice during the course of this work.*
- *Lastly but not leastly, I would like to thanks my families, both in law and bloodline, for their undying support and faith in me.*

Thanks for all the great times and helping me through. I couldn't have done it without you.



Bu-Ali Sina University

Faculty of Chemistry

In The Name of Allah

**Synthesis and characterization of macroacyclic and
macrocyclic complexes derived from various mono- and di-
aldehydes and some polyamines in the presence of different
metal ions**

**Thesis Submitted in Partial Fulfillment of the Requirements for the Degree of
Doctor of Philosophy (Ph.D.) in Inorganic Chemistry**

By:

Reza Azadbakht

Under The Supervision of:

Professor Hassan Keypour

Approved and evaluated by the thesis committee as:

H. Keypour, Prof. of Inorg. Chem. (Chairman).....*H...Keypour*.....

S. J. Sabounchei, Prof. of Inorg. Chem. (Internal Examiner)....*J..Sabounchei*.....

N. Safari, Prof. of Inorg. Chem. (External Examiner, Shahid Beheshti Univ.)...*N.Safari*.....

S. Amani, Prof. of Inorg. Chem. (External Examiner, Arak Univ.)....*Samiel.Amani*.....

M. A. Zolfigol, Prof. of Org. Chem. (Internal Examiner)....*M..A..Zolfigol*.....

S. Salehzadeh, Assoc. Prof. of Inorg. Chem. (Internal Examiner)*S..Salehzadeh*

February 2008

Abstract

Synthesis and characterization of macroacyclic and macrocyclic Complexes derived from various mono- and di- aldehydes and some polyamines in the presence of different metal Ions

By
Reza Azadbakht

In this project six new polyamine ligands and a number of novel complexes of corresponding macrocyclic and macro-acyclic Schiff-base ligands were synthesized and characterized.

Chapter 1: Introduction

Chapter 2: Polyamine ligands synthesis

Three new polyamine ligands of $N1-(2\text{-nitrobenzyl})-N1-(2\text{-aminoethyl})\text{ethane-}1,2\text{-diamine}$ (L^1), $N1-(2\text{-nitrobenzyl})-N1-(2\text{-aminoethyl})\text{propane-}1,3\text{-diamine}$ (L^2) and $N1-(2\text{-nitrobenzyl})-N1-(3\text{-aminopropyl})\text{propane-}1,3\text{-diamine}$ (L^3) were synthesized and their complexes with cadmium (II) ion were prepared. Three new asymmetric tripodal tetraamine ligands, $2\text{-}((\text{bis}(2\text{-aminoethyl})\text{amino})\text{methyl})\text{benzenamine}$ (L^4), $2\text{-}((2\text{-amino-ethyl})(3\text{-aminopropyl})\text{amino})\text{methyl}\text{benzenamine}$ (L^5) and $2\text{-}((\text{bis}(3\text{-aminopropyl})\text{amino-})\text{methyl})\text{benzenamine}$ (L^6) were synthesized by reducing of the only nitro group of the relevant L^1 , L^2 and L^3 ligands, so that leading to asymmetrical tripodal ligands that contain one primary aromatic amine group and two primary aliphatic amine groups.

Chapter 3: Synthesis and characterization of N_3O_2 macrocyclic Schiff-base complexes with 2-nitrobenzyl side chain

The cyclocondensation of three polyamine ligands of L^1 or L^2 and/or L^3 with $2\text{-}[2\text{-}(2\text{-formyl phenoxy)}\text{ethoxy]\text{benzaldehyde}}$ in the presence of various

metal(II) ions were examined. These reactions only in the presence of a stoichiometric amount of cadmium(II) ion gave related cadmium(II) macrocyclic Schiff-base complexes. In the other cases no cyclic complexes have been obtained and metal(II) polyamines were the only products. The complexes have been studied with IR, $^1\text{H}\text{NMR}$, $^{13}\text{C}\text{NMR}$, DEPT, COSY, HMQC and microanalysis. The crystal structures of $[\text{Cd}(\text{NO}_3)(L^7)(\mu-\text{NO}_3)\text{Cd}(\text{NO}_3)(L^7)]0.5\text{Cd}(\text{NO}_3)_4$ (1) (Fig. 3.1) and $[\text{Cd } L^7(\text{NO}_3)(\text{CH}_3\text{OH})]\text{ClO}_4$ (2) have been also determined.

Chapter 4: macrocyclic Schiff-base complexes with 2- aminobenzyl pendant arm

[1+1] Macrocyclic Schiff-base complexes containing 1, 2-diphenoxymethane head units and 2-aminobenzyl pendant arm, were synthesized as $[\text{MnL}^{10}(\text{MeOH})](\text{ClO}_4)_2$ (1), $[\text{MnL}^{11}(\text{MeOH})](\text{ClO}_4)_2$ (2), $[\text{MnL}^{12}(\text{MeOH})](\text{ClO}_4)_2$ (3), $[\text{CdL}^{10}(\text{H}_2\text{O})](\text{NO}_3)_2$ (4) and $[\text{CdL}^{11}(\text{H}_2\text{O})](\text{NO}_3)_2$ (5) from the metal ion templated cyclocondensation reactions of 2-[2-(2-formylphenoxy)ethoxy]benzaldehyde with (L^4), (L^5) or (L^6) tripodal tetraamine ligands. The crystal structure determination of (1) and (5) showed that the complex cations that had formed consisted of a pentagonal bipyramidally coordinated Mn(II) and Cd(II) ions, centrally located in a N_3O_2 macrocycle, with one 2-aminobenzyl pendant arm and a solvent molecule in axial positions. The cadmium complexes (4 and 5) have been studied by IR, $^1\text{H}\text{NMR}$, $^{13}\text{C}\text{NMR}$, DEPT, COSY, HMQC and microanalysis.

Chapter 5: Macroyclic ligand complexes containing pyridine moiety

Cyclocondensation of the asymmetric tripodal tetraamine ligand, 2-((bis(3-aminopropyl)amino)-methyl)benzenamine (L^6), in the presence of Ni(II), Cu(II) and Zn(II) ions, with 2,6-diacetylpyridine (DAP) were investigated. In the presence of Ni(II) ion, only a mono Schiff-base

macroacyclic complex was obtained but in the presence of Zn(II) and Cu(II) ions novel [14]-membered macrocyclic diimine Schiff-base complexes were the final products. Surprisingly, in both cases (Zn(II) and Cu(II)) metal ions 2-aminobenzyl unit is a part of macrocycle. The crystal structures of $[CuL^{14}](ClO_4)$ (2) and $[NiL^{15}](ClO_4)$ (3) were determined by X-ray diffraction. The zinc complex, $[ZnL^{14}](ClO_4)_2$ (1), has been studied by IR, 1H NMR, ^{13}C NMR, DEPT, COSY, HMQC and microanalysis.

Chapter 6: *Macroacyclic complexes.*

$[CdL^{17}(NO_3)_2]$, $[CdL^{18}(NO_3)_2]$, $[CdL^{19}(NO_3)_2]$, $[CoL^{17}(NO_3)_2]$, $[CoL^{18}(NO_3)_2]$, $[CoL^{19}(NO_3)_2]$, $[LaL^{20}]Cl_3$ and $[La_2L^{21}]Cl_2$ macroacyclic Schiff-base complexes were prepared by template condensation of 2-pyridinecarboxaldehyde, salicylaldehyde or 3-methoxysalicylaldehyde with L^1 or L and/or L^3 in the presence of corresponding metal ions. The Cd(II) and La(III) complexes have been studied with IR, 1H NMR, ^{13}C NMR, DEPT, COSY, HMQC and microanalysis. The complex $[CdL^{17}(NO_3)_2]$ (1) has been characterized through X-ray crystallography. In the complex 1 the Cd is in a six-coordinate environment comprised of the ligand N_4 -donor set and two oxygen atoms of two nitrate groups. In the case of reaction of 2-pyridinecarboxaldehyde and L_B (2.6.11.2) and Mn^{2+} one dication (9) compound was also synthesized and crystal structure of 9 has been determined with X-ray crystallography.

Chapter 7: *Computational studies*

As the crystal structure of some complexes could not be properly refined, we chose ab initio molecular orbital calculations to assess the structural parameters of these complexes in gas phase. We undertook a full geometry optimization at the LanL2MB, HF/3-21G and 6-31G* levels of theory, by using LanL2DZ basis set for some of metal ions. We also undertook the same level of ab initio calculations for complexes that we refine the crystal*

structures to observe the comparability of the predicted gas phase structural parameters with that determined in the solid state.

Table of contents

	page
Chapter 1: Introduction	
1.1. Background	2
1.2. Schiff-bases	5
1.3. Synthesis of Schiff-base Complexes	7
1.4. Ligand flexibility	9
1.5. Template ion size	10
1.6. Transmetallation reactions	14
References	16
Chapter 2: Polyamine Synthesis	
2.1. Introduction	23
2.2. Synthesis of tetraamine tripodal ligands	24
2.2.1. Tris(2-aminoethyl)amine	26
2.2.2. 2-Aminoethylbis(3-aminopropyl)amine (abap), Tris(3-aminopropyl)amine (trpn) and 4-Aminobutylbis(3-aminopropyl)amine (abbp)	26
2.3. Liner tetraamine ligands	35
2.4. Experimental	35
2.5. Safety note	36
2.6. Ligand synthesis	36
2.6.1. 3,3'-Diphthalimidodipropylamine	36
2.6.2. 2, 2'-Diphthalimidodiethylamine	36
2.6.3. 2, 3'-Diphthalimidoethylpropylamine	37
2.6.4. N1-(2-nitrobenzyl)-N1-(2-aminoethyl)ethane-1,2-diaminetrihydrochloride (I)	37
2.6.5. N1-(2-nitrobenzyl)-N1-(2-aminoethyl)propane-1,3-diaminetrihydrochloride (II)	39
2.6.5.1. N1-(2-nitrobenzyl)-N1-(2-aminoethyl)propane-1,3-diamine Threehydrochloride (II)	39
2.6.5.2. N1-(2-nitrobenzyl)-N1-(3-aminopropyl)propane-1,3-diamine Threehydrochloride (III)	40
2.6.6. 2-((bis(2-aminoethyl)amino)methyl)benzenamine (L^4)	40
2.6.7. 2-((2-aminoethyl)(3-aminopropyl)amino)methyl)benzenamine (L^5) and 2-((bis(3-aminopropyl)amino)methyl)benzenamine (L^6)	41
2.6.8. N,N'-bis(2-nitrophenyl)-1,2-ethandiamine(IV)	43

2.6.9.	N,N'-bis(2-nitrophenyl)-1,3-propanediamine(V) and N,N'-bis(2-nitrophenyl)-1,4-butanediamine(VI)	43
2.6.9.1.	N,N'-bis(2-nitrophenyl)-1,3-propanediamine(IV)	43
2.6.9.2.	N,N'-bis(2-nitrophenyl)-1,4-butanediamine(VI)	43
2.6.10.	N,N'-bis(2-aminophenyl)-1,2-ethanediamine(L ⁷)	43
2.6.11.	N,N'-bis(2-aminophenyl)-1,3-propanediamine(L ⁸) and N,N'-bis(2-aminophenyl)-1,4-butanediamine(L ⁹)	44
2.6.11.1	N,N'-bis(2-aminophenyl)-1,3-propanediamine(L ⁸)	44
2.6.11.2	N,N'-bis(2-aminophenyl)-1,4-butanediamine(L ⁹)	44
2.7.	Preparation of polyamine complexes	45
2.7. 1.	[CdL ¹ (NO ₃)]	45
2.7. 2.	[CdL ² (NO ₃) ₂]	46
2.7. 3.	[CdL ³ (NO ₃) ₂]	46
2.7.4.	[Ni (L ⁷) (CH ₃ CN) ₂](ClO ₄) ₂ , CH ₃ CN	46
2.8.	Result and discussion	46
2.8.1.	Synthesis and characterization	46
2.8.2.	X-ray crystal structure	47
2.9.	Conclusions	52
	References	52

Chapter 3: Synthesis and characterization of Schiff-base macrocyclic N₃O₂ complexes with 2-nitrobenzylpendant

3.1.	Introduction	62
3.2.	Experimental	71
3.3.	Starting materials	71
3.4.	Instrumentation	71
3.5.	X-ray crystal structure determination	72
3.6.	Synthesis	73
3.6.1.	2-[2-(2-formyl phenoxy)ethoxy]benzaldehyde	73
3.6.2.	Preparation of complexes	74
3.6.2.1.	General synthesis	74
3.6.2.2.	[Cd(NO ₃) (L ₅) (μ-NO ₃) Cd(NO ₃) (L ₅)] 0.5Cd(NO ₃) ₄ or [Cd(L ₅)] ²⁺	74
3.6.2.3.	[Cd(NO ₃)(CH ₃ OH) L ₅]ClO ₄ .C ₂ H ₆ O	74

3.6.2.4.	$[\text{CdL}_6](\text{NO}_3)_2$ (3)	75
3.6.2.5.	$[\text{CdL}_7](\text{NO}_3)_2$ (4)	75
3.7.	Results and discussion	76
3.7.1.	Synthesis and characterization	76
3.7.2.	X-ray crystal structure	77
3.8.	NMR study	81
3.9.	Conclusion	84
3.10.	Supplementary material	85
	References	85

Chapter 4: Macrocyclic complexes with 2-aminobenzyl pendant arm

4.1.	Introduction	93
4.2.	Experimental	102
4.2.1.	Starting materials	102
4.2.2.	Instrumentation	103
4.2.3.	X-ray crystal structure determination	103
4.2.4.	Preparation of complexes	104
4.2.4.1.	General procedure	104
4.2.4.2.	$[\text{MnL}^{10}(\text{CH}_3\text{OH})](\text{ClO}_4)_2$	105
4.2.4.3.	$[\text{MnL}^{11}(\text{CH}_3\text{OH})](\text{ClO}_4)_2$	105
4.2.4.4.	$[\text{MnL}^{12}(\text{CH}_3\text{OH})](\text{ClO}_4)_2$	105
4.2.4.5.	$[\text{CdL}^{10}(\text{H}_2\text{O})](\text{NO}_3)_2$	106
4.2.4.6.	$[\text{CdL}^{11}(\text{H}_2\text{O})](\text{NO}_3)_2$	106
4.3.	Results and Discussion	107
4.3.1.	Synthesis and characterization	107
4.3.2.	X-ray crystal structure	107
4.4.	NMR Studies	114
4.5.	$[\text{MnL}^{13}]\text{ClO}_4$	116
4.5.	Conclusions	117
4.6.	Supplementary material	117
	References	118

Chapter 5: Macrocyclic ligand complexes containing pyridine moiety

5.1.	Introduction	122
5.2.	Macrocyclic ligand complexes containing pyridine moiety	122

5.3.	Macrocycles with pendant functional group	141
5.4.	Experimental	149
5.4.1.	Preparation of complexes	151
5.4.1.1.	$[\text{ZnL}^{14}](\text{ClO}_4)_2$ (1)	151
5.4.1.2.	$[\text{CuL}^{14}](\text{ClO}_4)_2 \cdot \text{CH}_3\text{OH}$ (2)	151
5.4.1.3.	$[\text{NiL}^{15}](\text{ClO}_4)_2 \cdot \text{CH}_3\text{CN}$ (3)	151
5.5.	Result and discussion	152
5.5.1.	Synthesis of Complexes.	152
5.5.2.	X-ray crystal structure	154
5.6.	NMR studies	159
5.7.	$[\text{NiL}^{16}]\text{ClO}_4$	160
5.7.	Conclusion	161
	Refrences	162

Chapter 6: Macroacyclic complexes

6.1.	Introduction	174
6.2.	Experimental	182
6.2.1.	Starting materials	182
6.2.2.	Instrumentation	182
6.2.3.	X-ray crystal structure determination	183
6.3.	Synthesis	184
6.3.1.	Preparation of macroacyclic complexes of Cd(II) and Co(II)	184
6.3.2.	General synthesis	184
6.3.3.	$[\text{Cd L}^{17}(\text{NO}_3)_2]$ (1)	185
6.3.4.	$[\text{CdL}^{18}(\text{NO}_3)_2]$ (2)	185
6.3.5.	$[\text{CdL}^{19}(\text{NO}_3)_2]$ (3)	186
4.3.6.	$[\text{CoL}^{17}(\text{NO}_3)_2]$ (4)	186
6.3.7.	$[\text{Co L}^{18}(\text{NO}_3)_2]$ (5)	186
6.3.8.	$[\text{CoL}^{19}(\text{NO}_3)_2]$ (6)	186
6.3.9.	Preparation of macroacyclic complexes of La(III)	189
6.3.9.1.	$[\text{La L}^7] \text{Cl}_3$ (7)	189
6.3.9.2.	$[\text{La}_2 \text{L}^8] \text{Cl}_2$ (8)	189

6.4.	Results and discussion	190
6.4.1.	Synthesis and characterization	190
6.4.2.	X-ray crystal structure	191
6.5.	NMR Studies	192
6.6.	Synthesis carbocation	196
6.7.	Results and discussion	198
6.8.	Supplementary material	198
	References	198

Chapter 7: Computational studies

7-1.	Introduction	205
7-2.	Basis Set	208
7-2-1.	Minimal Basis Sets	208
7-2-2.	Split Valance Basis Sets	209
7-2-3.	Polarized Basis Sets	209
7-2-4.	Basis Sets for Post-Third-Row Atoms	210
7-3.	ab initio modelling studies of Ni(II) Polyamine Complex	210
7.4.	macrocyclic (ab initio modelling studies of Cd(II) and Mn(II) Complexes	211
7.5.	Ab initio modelling studies of Cd(II) macroacyclic Complexes	215
	References	217

List of table

Table 2. 1.	Crystal data and structure refinement for $[\text{NiL}_B(\text{CH}_3\text{CN})_2](\text{ClO}_4)_2$ (4) and $[\text{CdL}^1(\text{NO}_3)_2]$ (1)	48
Table 2.2.	Important bond lengths (\AA) and angles ($^\circ$) for $[\text{Ni}(\text{L}^7)(\text{CH}_3\text{CN})_2]^{2+}$ and $[\text{CdL}^1(\text{NO}_3)_2]$	50
Table 2.3.	Torsion angles ($^\circ$) for $[\text{CdL}^1(\text{NO}_3)_2]$ (1)	50
Table 2.4.	Torsion angles ($^\circ$) for $[\text{Ni L}^7 (\text{CH}_3\text{CN})_2](\text{ClO}_4)_2$ (4)	51
Table 3. 1.	Table 3. 1. Crystal data and structure refinement for $[\text{Cd}(\text{NO}_3)(\text{L}^7)(\mu\text{-NO}_3)\text{Cd}(\text{NO}_3)(\text{L}^7)] \cdot 0.5\text{cd}(\text{NO}_3)_4$ (1) and $[\text{Cd L}^7(\text{NO}_3)(\text{CH}_3\text{OH})]\text{ClO}_4 \cdot \text{C}_2\text{H}_6\text{O}$ (2)	72
Table 3. 2.	Important bond lengths (\AA) and Bond angle ($^\circ$) for $[\text{Cd}(\text{NO}_3)(\text{CH}_3\text{OH})\text{L}^7]\text{ClO}_4$	79
Table 3. 3.	Torsion angles for $[\text{Cd}(\text{NO}_3)(\text{CH}_3\text{OH})\text{L}^7]^{2+}$	80
Table 3. 4.	^1H NMR and ^{13}C NMR {1H} spectral assignments for $[\text{CdL}^7]^{2+}$, $[\text{CdL}^8]^{2+}$, $[\text{CdL}^9]^{2+}$, recorded in DMSO-d6	82
Table 4. 1.	Crystal data and structure refinement for $[\text{MnL}^{10}(\text{CH}_3\text{OH})](\text{ClO}_4)_2$ (1) and $[\text{CdL}^{11}(\text{H}_2\text{O})](\text{NO}_3)_2$ (4).	104
Table 4. 2.	Selected bond lengths (\AA) and bond angles ($^\circ$) for $[\text{MnL}^{10}(\text{CH}_3\text{OH})]^{2+}$ and $[\text{Cd L}^{11}(\text{H}_2\text{O})]^{2+}$	108
Table 4. 3.	Torsion angles of $[\text{MnL}^{10}(\text{CH}_3\text{OH})]^{2+}$	111
Table 4. 4.	Other bond angles for $[\text{CdL}^{11}(\text{H}_2\text{O})]^{2+}$	129
Table 4. 5.	^1H NMR and ^{13}C NMR {1H} spectral assignments for $[\text{CdL}^{10}]^{2+}$ and $[\text{CdL}^{11}]^{2+}$, recorded in DMSO.	132
Table 5. 1.	Crystal data and structure refinement for $[\text{NiL}^{15}](\text{ClO}_4)_2$ and $[\text{CuL}^{14}](\text{ClO}_4)_2 \cdot \text{CH}_3\text{OH}$	141
Table 5. 2.	Important bond lengths (\AA) and Bond angle ($^\circ$) $[\text{NiL}^{15}](\text{ClO}_4)_2 \cdot \text{CH}_3\text{CN}$ (3) and $[\text{CuL}^{14}(\text{H}_2\text{O})](\text{ClO}_4)_2 \cdot \text{CH}_3\text{OH}$ (2)	157
Table 5. 3.	Torsion angle ($^\circ$) of $[\text{NiL}^{15}](\text{ClO}_4)_2 \cdot \text{CH}_3\text{CN}$	157
Table 5. 4.	Torsion angles ($^\circ$) of $[\text{CuL}^{14}(\text{H}_2\text{O})]^{2+}$	158
Table 5. 5.	^1H NMR and ^{13}C NMR {1H} spectra assignments for $[\text{ZnL}^{14}](\text{ClO}_4)_2$ recorded in (DMSO-d6)	160

Table 6. 1.	Crystal data and structure refinement for $[\text{CdL}^{17}(\text{NO}_3)_2]$ and $\text{C}_{38}\text{H}_{32}\text{N}_8(\text{ClO}_4)_2$	183
Table 6. 2.	Selected bond lengths (\AA) and angles ($^\circ$) for $[\text{CdL}^{17}(\text{NO}_3)_2]$	187
Table 6. 3.	Other parameters ($^\circ$) for $[\text{CdL}^{17}(\text{NO}_3)_2]$	187
Table 6. 4.	^1H NMR and ^{13}C NMR{ ^1H } spectral assignments for 1, 2, 3 and related polyamine complex recorded in DMSO	194
Table 6.5.	Selected bond lengths (\AA) and angles ($^\circ$) for $\text{C}_{38}\text{H}_{32}\text{N}_8(\text{ClO}_4)_2$	198
Table 7. 1.	Selected bond lengths (\AA) and angles ($^\circ$) for $[\text{NiL}_B(\text{CH}_3\text{CN})_2]^{2+}$	211
Table 7. 2.	Selected bond lengths (\AA) and angles ($^\circ$) for $[\text{MnL}^{10}(\text{CH}_3\text{OH})]^{2+}$ (1), $[\text{MnL}^{11}(\text{CH}_3\text{OH})]^{2+}$ (2), $[\text{MnL}^{12}(\text{CH}_3\text{OH})]^{2+}$ (3), $[\text{CdL}^{10}(\text{H}_2\text{O})]^{2+}$ (4), $[\text{CdL}^{11}(\text{H}_2\text{O})]^{2+}$ (5) and $[\text{CdL}^{12}(\text{H}_2\text{O})]^{2+}$ (6)	214
Table 7. 3.	The ab initio optimized structures of the $[\text{Cd}(\text{L}^{17})(\text{NO}_3)_2]$ (1), $[\text{Cd}(\text{L}^{18})(\text{NO}_3)_2]$ (2) and $[\text{Cd}(\text{L}^{19})(\text{NO}_3)_2]$ (3)	216

List of figures

Fig. 1.1.	ORTEP drawing of the $[Ni(4)]^{2+}$ cation and related macrocycle.	12
Fig. 2.1.	General structur of tetradentate tripododal ligand.	23
Fig. 2.2.	skeleton types of liner tetraamines.	35
Fig. 2.3.	Structure of $[CdL^1(NO_3)_2]$ (1) and $[NiL_B(CH_3CN)_2](ClO_4)_2$ (4)	49
Fig. 3.1.	Structure of the cation $[Cd(NO_3)(L^7)(\mu\text{-}NO_3) Cd (NO_3)(L^7)]^+$	79
Fig. 3.2.	ORTEP depiction of $[Cd(NO_3)(CH_3OH)L^7]^+$ (2) and similar complex (5)	80
Fig. 4.1.	Schiff-base macrocycles L^1 - L^6	95
Fig. 4.2.	Structures of the cations $[MnL^{10}(CH_3OH)]^{2+}$ and $[CdL^{11}(H_2O)]^{2+}$	110
Fig. 5.1.	Structures of the cations $[CuL^{14}(H_2O)]^{2+}$ and $[NiL^{15}]^{2+}$	156
Fig. 6.1.	ORTEP drawing of the molecular structures of the complexes cation from $[CdL^{17}(NO_3)_2]$	187
Fig. 6.2 .	ORTEP drawing of the molecular structures of the dication, $C_{38}H_{32}N_8(ClO_4)_2$	197
Fig. 7.1.	The ab initio optimized structure of the cation $[NiL_B(CH_3CN)_2]^{2+}$.	210
Fig. 7.2.	The ab initio optimized structhrs of the cations $[MnL^{10}(CH_3OH)]^{2+}$, $[MnL^{11}(CH_3OH)]^{2+}$, $[MnL^{12}(CH_3OH)]^{2+}$, $[CdL^{10}(H_2O)]^{2+}$, $[CdL^{11}(H_2O)]^{2+}$ and $[CdL^{12}(H_2O)]^{2+}$	213
Fig. 7.3.	The ab initio optimized structures of the $[Cd(L^{17})(NO_3)_2]$ (1), $[Cd(L^{18})(NO_3)_2]$ (2) and $[Cd(L^{19})(NO_3)_2]$ (3)	216

List of Spectra

Spec. No. 2.1.	Infrared spectrum of N1-(2-nitrobenzyl)-N1-(2-aminoethyl)ethane-1,2-diaminetrihydrochloride (I) recorded as KBr pellet	219
Spec. No. 2.2.	¹ H NMR spectrum of N1-(2-nitrobenzyl)-N1-(2-aminoethyl)ethane-1,2-diaminetrihydrochloride (I)	220
Spec. No. 2.3.	¹³ C NMR spectrum of N1-(2-nitrobenzyl)-N1-(2-aminoethyl)ethane-1,2-diaminetrihydrochloride (I)	221
Spec. No. 2.4.	Infrared spectrum of N1-(2-nitrobenzyl)-N1-(2-aminoethyl)propane-1,3-diaminetrihydrochloride (II) recorded as KBr pellet	222
Spec. No. 2.5.	¹ H NMR spectrum of N1-(2-nitrobenzyl)-N1-(2-aminoethyl)propane-1,3-diaminetrihydrochloride (II)	223
Spec. No. 2.6.	¹³ C NMR spectrum of N1-(2-nitrobenzyl)-N1-(2-aminoethyl)propane-1,3-diaminetrihydrochloride (II)	224
Spec. No. 2.7.	¹ H NMR spectrum of N1-(2-nitrobenzyl)-N1-(3-aminopropyl)propane-1,3-diaminetrihydrochloride (III)	225
Spec. No. 2.8.	¹³ C NMR spectrum of N1-(2-nitrobenzyl)-N1-(3-aminopropyl)propane-1,3-diaminetrihydrochloride (III)	226
Spec. No. 2.9.	¹ H NMR spectrum of 2-((bis(2-aminoethyl)amino)methyl)benzenamine (<i>L</i> ⁴)	227
Spec. No. 2.10.	¹³ C NMR spectrum of 2-((bis(2-aminoethyl)amino)methyl)benzenamine (<i>L</i> ⁴)	228
Spec. No. 2.11.	¹ H NMR spectrum of 2-((bis(2-aminoethyl)amino)methyl)benzenamine (<i>L</i> ⁴)+D ₂ O	229
Spec. No. 2.12.	¹ H NMR spectrum of 2-((2-aminoethyl)(3-aminopropyl)amino)methyl)benzenamine (<i>L</i> ⁵)	230
Spec. No. 2.13.	¹³ C NMR spectrum of 2-((2-aminoethyl)(3-aminopropyl)amino)methyl)benzenamine (<i>L</i> ⁵)	231
Spec. No. 2.14.	¹ H NMR spectrum of 2-((bis(3-aminopropyl)amino)methyl)benzenamine (<i>L</i> ⁶)	232
Spec. No. 2.15.	¹ H NMR spectrum of 2-((bis(3-aminopropyl)amino)methyl)benzenamine (<i>L</i> ⁶)+ D ₂ O	233
Spec. No. 2.16.	¹³ C NMR spectrum of 2-((bis(3-aminopropyl)amino)methyl)benzenamine (<i>L</i> ⁶)	234
Spec. No. 2.17.	Infrared spectrum of [CdL ¹ (NO ₃)] (1) recorded as KBr pellet	235
Spec. No. 2.18.	¹ H NMR spectrum of [CdL ¹ (NO ₃)] (1)	236
Spec. No. 2.19.	¹³ C NMR spectrum of [CdL ¹ (NO ₃)] (1)	237
Spec. No. 2.20.	Infrared spectrum of [CdL ² (NO ₃)] (2) recorded as KBr pellet	238
Spec. No. 2.21.	¹ H NMR spectrum of [CdL ² (NO ₃)] (2)	239
Spec. No. 2.22.	¹³ C NMR spectrum of [CdL ² (NO ₃)] (2)	240